

IDS #7

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/63, 15/82	A1	(11) International Publication Number: WO 98/53083 (43) International Publication Date: 26 November 1998 (26.11.98)
--	----	--

(21) International Application Number: PCT/GB98/01450

(22) International Filing Date: 20 May 1998 (20.05.98)

(30) Priority Data:  
9710475.6 (21 May 1997) (21.05.97) GB

(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GRIERSON, Donald [GB/GB]; University of Nottingham, Sutton Bonnington Campus, Loughborough LE12 5RD (GB). LOWE, Alexandra, Louise [GB/GB]; University of Nottingham, Sutton Bonnington Campus, Loughborough LE12 5RD (GB). HAMILTON, Andrew, John [GB/GB]; University of Nottingham, Sutton Bonnington Campus, Loughborough LE12 5RD (GB).

(74) Agents: HUSKISSON, Frank, Mackie et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG42 6YA (GB).


(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With International search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: GENE SILENCING



(57) Abstract

Constructs and methods for enhancing the inhibition of a target gene within an organism involve inserting into the gene silencing vector an inverted repeat sequence of all or part of a polynucleotide region within the vector. The inverted repeat sequence may be a synthetic polynucleotide sequence or comprise a modified natural polynucleotide sequence.

Post-transcriptional?  
What is the phenotype?  
Protein & RNA?

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	ME	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

## GENE SILENCING

This invention relates to the control of gene expression, more particularly to the inhibition of expression, commonly referred to as "gene silencing".

5

Two principal methods for the modulation of gene expression are known. These are referred to in the art as "antisense downregulation" and "sense downregulation" (also, referred to as "cosuppression"). Both of these methods lead to an inhibition of expression of the target gene.

10

In antisense downregulation, a DNA which is complementary to all or part of an endogenous target gene is inserted into the genome in reverse orientation. While the mechanism has not been fully elucidated, one theory is that transcription of such an antisense gene produces mRNA which is complementary in sequence to the mRNA product transcribed from the endogenous gene: that antisense mRNA then binds with the naturally produced "sense" mRNA to form a duplex which inhibits translation of the natural mRNA to protein. It is not necessary that the inserted antisense gene be equal in length to the endogenous gene sequence: a fragment is sufficient. The size of the fragment does not appear to be particularly important. Fragments as small as 42 or so nucleotides have been reported to be effective. Generally somewhere in the region of 50 nucleotides is accepted as sufficient to obtain the inhibitory effect. However, it has to be said that fewer nucleotides may very well work: a greater number, up to the equivalent of full length, will certainly work. It is usual simply to use a fragment length for which there is a convenient restriction enzyme cleavage site somewhere downstream of fifty nucleotides. The fact that only a fragment of the gene is required means that not all of the gene need be sequenced. It also means that commonly a cDNA will suffice, obviating the need to isolate the full genomic sequence.

15

20

25

30

The antisense fragment does not have to be precisely the same as the endogenous complementary strand of the target gene. There simply has to be sufficient sequence similarity to achieve inhibition of the target gene. This is an important feature of antisense technology as it permits the use of a sequence which has been derived from one plant species to be effective in another and obviates the need to construct antisense vectors for each

- 2 -

individual species of interest. Although sequences isolated from one species may be effective in another, it is not infrequent to find exceptions where the degree of sequence similarity between one species and the other is insufficient for the effect to be obtained. In such cases, it may be necessary to isolate the species-specific homologue.

5 Antisense downregulation technology is well-established in the art. It is the subject of several textbooks and many hundreds of journal publications. The principal patent reference is European Patent No. 240,208 in the name of Calgene Inc. There is no reason to doubt the operability of antisense technology. It is well-established, used routinely in laboratories around the world and products in which it is used are on the market.

10 Both overexpression and downregulation are achieved by "sense" technology. If a full length copy of the target gene is inserted into the genome then a range of phenotypes is obtained, some overexpressing the target gene, some underexpressing. A population of plants produced by this method may then be screened and individual phenotypes isolated. A similarity with antisense is that the inserted sequence need not be a full length copy. The principal patent reference on cosuppression is European Patent 465,572 in the name of DNA  
15 Plant Technology Inc. There is no reason to doubt the operability of sense/cosuppression technology. It is well-established, used routinely in laboratories around the world and products in which it is used are on the market.

Sense and antisense gene regulation is reviewed by Bird and Ray in Biotechnology  
20 and Genetic Engineering Reviews 9: 207-227 (1991). The use of these techniques to control selected genes in tomato has been described by Gray et.al., Plant Molecular Biology, 19: 69-87 (1992).

Gene silencing can therefore be achieved by inserting into the genome of a target organism an extra copy of the target gene coding sequence which may comprise either the  
25 whole or part or be a truncated sequence and may be in sense or antisense orientation. Additionally, intron sequences which are obtainable from the genomic gene sequence may be used in the construction of suppression vectors. There have also been reports of gene silencing being achieved within organisms of both the transgene and the endogenous gene where the only sequence identity is within the promoter regions.

30

- 3 -

Gene control by any of the methods described requires insertion of the sense or antisense sequence, under control of appropriate promoters and termination sequences containing polyadenylation signals, into the genome of the target plant species by transformation, followed by regeneration of the transformants into whole plants. It is probably fair to say  
5 that transformation methods exist for most plant species or can be obtained by adaptation of available methods.

The most widely used method is *Agrobacterium*- mediated transformation, mainly for dicotyledonous species. This is the best known, most widely studied and, therefore, best understood of all transformation methods. The rhizobacterium *Agrobacterium tumefaciens*,  
10 or the related *Agrobacterium rhizogenes*, contain certain plasmids which, in nature, cause the formation of disease symptoms, crown gall or hairy root tumours, in plants which are infected by the bacterium. Part of the mechanism employed by *Agrobacterium* in pathogenesis is that a section of plasmid DNA which is bounded by right and left border regions is transferred stably into the genome of the infected plant. Therefore, if foreign DNA  
15 is inserted into the so-called "transfer" region (T-region) in substitution for the genes normally present therein, that foreign gene will be transferred into the plant genome. There are many hundreds of references in the journal literature, in textbooks and in patents and the methodology is well-established.

Various methods for the direct insertion of DNA into the nucleus of monocot cells  
20 are known.

In the ballistic method, microparticles of dense material, usually gold or tungsten, are fired at high velocity at the target cells where they penetrate the cells, opening an aperture in the cell wall through which DNA may enter. The DNA may be coated on to the microparticles or may be added to the culture medium.

25 In microinjection, the DNA is inserted by injection into individual cells via an ultrafine hollow needle.

Another method, applicable to both monocots and dicots, involves creating a suspension of the target cells in a liquid, adding microscopic needle-like material, such as silicon carbide or silicon nitride "whiskers", and agitating so that the cells and whiskers  
30 collide and DNA present in the liquid enters the cell.

- 4 -

In summary, then, the requirements for gene silencing using both sense and antisense technology are known and the methods by which the required sequences may be introduced are known.

The present invention aims to, *inter alia*, provide a method of enhancing the control  
5 of gene expression.

According to the present invention there is provided a vector for enhancing the inhibition of a selected target gene within an organism, comprising a gene silencing vector characterised in that the said gene silencing vector includes a inverted repeat of all or part of  
a polynucleotide region within the vector. ||

10 ~~The inverted repeat sequence may be a synthetic polynucleotide sequence and its inverted repeat sequence or an inverted repeat of all or part of the said gene silencing vector or an inverted repeat of the 5'-untranslated region of the gene silencing vector.~~

The inverted repeat may be separated from the polynucleotide region by a sequence of nucleotides.

15 The invention also provides a method of controlling the expression of a DNA sequence in a target organism, comprising inserting into the genome of said organism an enhanced gene silencing vector as defined above.

In a preferred embodiment a vector for enhanced gene silencing comprising in sequence a promoter region, a 5'-untranslated region, a transcribable DNA sequence and a  
20 3'-untranslated region containing a polyadenylation signal, characterised in that the said construct includes an inverted repeat of a region of said vector.

~~It is preferred that the inverted repeat is a fragment of the 5'-untranslated region of the said vector. The vector may have two tandem copies of the inverted repeat.~~

In simple terms, we have found that the inhibitory effect of a gene-silencing vector  
25 can be enhanced by creating in the vector an inverted repeat of a part of the sequence of the vector. Alternatively the inverted repeat may be of a synthetic sequence which may be created independently of the vector itself and then inserted into the vector sequence. While the mechanism by which the enhancement is achieved is not fully understood we understand that the minimum required for such a vector is a region or regions which identify the gene  
30 targeted for silencing and an inverted repeat of a part of that region or, as explained above an inserted sequence and its inverted repeat. The region of the vector which identifies the gene

- 5 -

targeted for silencing may be any part of that endogenous gene which characterises it, for example, its promoter, its 5'-untranslated region, its coding sequence or its 3'untranslated region. ~~We have also found that the vector used in this invention will silence the expression of the target gene and also any members of the gene family to which the targeted gene~~  
5 belongs.

Although the mechanism by which the invention operates is not fully understood, we believe that creation of an inverted repeat promotes the formation of a duplex DNA between the selected sequence and its inverted.

The inverted repeat may be positioned anywhere within the vector such as within the  
10 promoter region, the 5' untranslated region, the coding sequence or the 3' untranslated region. If the inverted repeat is based on a contiguous sequence within the promoter region, then it is preferred that the inverted repeat is located within the promoter region. If the inverted repeat is based on a contiguous sequence within the 5' untranslated region, then it is preferred that the inverted repeat is located within the 5' untranslated region. ~~If the inverted~~  
15 ~~repeat is based on a contiguous sequence within the coding region, then it is preferred that~~  
~~the inverted repeat is located within the coding region.~~ If the inverted repeat is based on a contiguous sequence within the 3' untranslated region, then it is preferred that the inverted repeat is located within the 3' untranslated region.

The selected polynucleotide sequence and its inverted repeat may or may not be  
20 separated by a polynucleotide sequence which remains unpaired when the 5' untranslated region and the inverted repeat have formed a DNA duplex. It is preferred however, that the chosen contiguous sequence and its inverted repeat are separated by a polynucleotide sequence which remains unpaired when the 5' untranslated region and the inverted repeat have formed a DNA duplex.

25 It is further preferred that the inverted repeat is based on the 5' untranslated sequence. It is also preferred that the inverted repeat is positioned upstream of the coding sequence. It is further preferred that the inverted repeat is positioned between the 5' untranslated region and the coding sequence. It is further preferred that the 5' untranslated region and the inverted repeat are separated by a polynucleotide sequence which remains unpaired when the  
30 5' untranslated region and the inverted repeat have formed a DNA duplex.

- 6 -

Suppression can also be achieved by creating a vector containing an inverted repeat sequence which is capable of forming a duplex DNA within the promoter region of the target gene. This obviates the need to include any specific coding sequence information about the gene to be suppressed since the vector would allow suppression of the promoter within the organism and hence the expression of the target gene. Alternatively vectors may be created which are lacking a promoter sequence but which contain an inverted repeat of a sequence within the 5' untranslated region, the coding region or the 3' untranslated region.

The 5' or 3' untranslated regions of a gene suppression vector can also be replaced with a synthetic 5' or 3' untranslated regions which comprises a polynucleotide part and inverted repeat separated by a polynucleotide sequence which remains unpaired when the said polynucleotide part and the inverted repeat form a DNA duplex. It is preferred to construct a synthetic 5' untranslated region. It is further preferred to construct the synthetic 5' untranslated region comprising sequentially, a 33 base polynucleotide part and a 33 base polynucleotide inverted repeat separated by a 12 base polynucleotide.

Where it is desired to use an inverted repeat sequence within the 5' untranslated region, the coding sequence or the 3' untranslated region, gene silencing vectors constructed with inverted repeats within any one of these regions may additionally enable the silencing of genes that are homologous to the coding sequence present in the silencing vector. Therefore when it is desired to silence genes homologues within an organism the construction of a silencing vector containing an inverted repeat within the 5' untranslated region, the coding sequence or the 3' untranslated region may allow the silencing of all the genes exhibiting sequence homology to the coding sequence within the construct. Homology/homologous usually denotes those sequences which are of some common ancestral structure and exhibit a high degree of sequence similarity of the active regions. Examples of homologous genes include the ACC-oxidase enzyme gene family which includes ACO1 and ACO2.

Any of the sequences of the present invention may be produced and manipulated using standard molecular biology techniques. The sequences may be obtained from a desired organism source such as plant sources and modified as required or synthesised *ab initio* using standard oligosynthetic techniques.



- 7 -

Without wishing to be bound by any particular theory of how it may work, the following is a discussion of our invention. 96% of tomato plants transformed with an ACC-oxidase sense gene containing two additional, upstream inverted copies of its 5' untranslated region, exhibited substantially reduced ACC-oxidase activity compared to wild type plants. Only 15% of plants transformed with a similar construct, without the inverted repeat, had reduced ACC-oxidase activity. Both populations had similar average numbers of transgenes per plant. Treatment of tomato leaves with cycloheximide caused a strong, reproducible increase in the abundance of ACC-oxidase transcripts and was used in the study of suppression by ACC-oxidase sense transgenes in preference to wound induction used in previous studies. The relative abundance of unprocessed and processed ACC-oxidase transcripts in suppressed and non-suppressed plants was assayed by ribonuclease protection assays, providing an indirect measure of transcription and mRNA accumulation which did not rely upon assaying isolated nuclei. This analysis indicated that the suppression of ACO1 gene expression was mainly post-transcriptional. Using the same type of RPA assay similar results were obtained from plants containing suppressing polygalactonase-sense or ACO-antisense transgenes.

There are now numerous examples of the inactivation of homologous sequences in plants. The term "homology dependent gene silencing" (HDGS) best describes all of these although it should be noted that in most examples the "silencing" is not complete and a low level of gene expression remains. Throughout this specification we will use the classification most-recently outlined by Matzke and Matzke, Plant Physiol. 107: 679-685 (1995) in which different examples of HDGS were divided into three main groups; cis-inactivation, trans-inactivation, and sense-suppression. Down regulation by antisense genes bears many similarities to the last of these and has been suggested to operated by the same mechanism (Grierson et al, Trends Biotechnol. 9: 122-123 (1991)). Both sense and antisense transgenes have been widely used to reduce the expression of homologous endogenous genes in plants. Although the underlying mechanisms of HDGS remain obscure, this technology has found numerous applications not only in fundamental research but also in commercial biotechnology ventures and new food products are already on the market.

At present, obtaining a large number of strongly suppressed, transgenic lines is more a matter of luck than judgement. A positive correlation between the presence of repeated

- 8 -

transgene sequences and the incidence of HDGS has been noted. However single locus-transgene insertions associated with HDGS have also been reported.

There is an emerging consensus that different examples of HDGS can be classified on the basis of whether or not the transcription of the target gene is affected. Examples of transcriptional suppression have been described. Where the homology between interacting genes resides within transcribed sequences, HDGS has been shown to be a post-transcriptional effect. Despite this apparently precise demarcation, several similarities exist between some examples in the two different categories. These include variegated patterns of silencing, increased methylation of genes participating in silencing and the frequent observation that silencing loci contain repeated sequences.

Although transcriptional silencing must occur in the nucleus, post-transcriptional silencing might occur in either or both the nucleus or cytoplasm. There is evidence that the abundance of processed, nuclear RNA of silenced genes was unaffected and suggested an effect upon transport into or degradation within the cytoplasm. More compelling evidence that post-transcriptional HDGS occurs outside the nucleus is the relationship between gene silencing involving nuclear transgenes and resistance to cytoplasmically replicating RNA viruses. Transgenic plants containing transgenes that suppress the activity of other transgenes (e.g. GUS) or endogenous genes (e.g. PG) are also resistant to RNA viruses which have been engineered to include sequences from those genes. Nevertheless, nuclear features such as transgene methylation and complexity of transgene loci were found positively to correlate with virus resistance. In almost all instances of HDGS, the source of the silencing is nuclear (even if the manifestation is cytoplasmic). However, silencing of a nuclear gene by a cytoplasmic element has been demonstrated by the suppression of phytoene desaturase in plants infected by a recombinant virus containing sequences from that gene.

Although, there are now numerous examples of post-transcriptional suppression of plant genes by HDGS, as yet, there is no information as to whether the increased turnover of pre-mRNA is related to or distinct from other cellular, RNA turnover processes. Degradation of RNA in plants is poorly understood but there is evidence that translation is involved. For example, the very short half lives (around 10 minutes) of small auxin up RNAs (SAURS) can be markedly prolonged by treatment with cycloheximide.

- 9 -

This invention gives a striking increase in the frequency of HDGS following the inclusion of a short repeated region within a transgene. Expression of the target gene encoding the terminal ethylene biosynthetic enzyme ACC-oxidase, in tomato was suppressed by such constructs mainly post-transcriptionally. This was shown to be true for other  
5 examples of sense and antisense suppression in tomato. Cycloheximide was found to be a potent and reliable inducer of ACO gene expression but did not ameliorate the silencing.

The invention will now be described, by way of illustration, in the following Examples and with reference to the accompanying Figures of which:

10 **Figure 1.** (A) ACO1 gene silencing vector.

(B) ACO1 gene silencing vector containing tandem inverted repeats of the 5' untranslated region.

**Figure 2.** Illustrates the relative ACC-oxidase activity in both types of transgenic plant relative to wild type values where C = transgenic plants containing construct C (Figure 1A)  
15 and V = transgenic plants containing construct V (Figure 1B).

**Figure 3.** Tomato plant ACC Oxidase activity of transgenic transformants containing  
[ pHIR-ACO (as illustrated in SEQ ID No 10). The graph also includes C12ACO (overexpression control) an untransformed wild type and TOM13 strong antisense gene silenced control.

## 20 **Example 1.0**

Construct V (Figure 1) was made in the following manner: 79 base pairs of the 5' untranslated region of the tomato ACO1 cDNA was amplified by PCR and two copies were ligated in tandem in the reverse orientation immediately upstream of the ACO1 cDNA which contains its own polyadenylation signal in its 3' untranslated region (construct C). Both  
25 were ligated downstream of the CaMV 35S promoter and then transferred to the binary vector, Bin19. Figure 1 shows the basic details of constructs "C" and "V". These were used to transform tomato plants (Ailsa Craig) by *Agrobacterium* mediated DNA transfer. 13 and

- 10 -

28 individual kanamycin resistant calli were obtained with constructs "C" and "V" respectively and these were regenerated into plants.

The nucleotide sequence of the promoter and 5' untranslated region of the ACO1 gene is given as SEQ ID NO 1 hereinafter. The 79bp referred to above begins at base  
5 number 1874 and stops at the base immediately preceding the translation start codon (ATG) at number 1952.

#### Example 1.1

To screen the population for any effects on ACO gene expression, relative ACO activity was measured from untransformed and transformed plants. The production of  
10 ethylene from leaf discs supplemented with the ethylene precursor, 1-aminocyclopropane-1-carboxylic acid, was measured at least three times from each plant. The cutting of the discs by a cork borer wounds the leaves and stimulates the expression of the ACO1 gene. ACC-oxidase activity in both types of transgenic plant relative to wild type values are shown in Figure 2. There was a dramatic difference in ACO activity between the two populations,  
15 with plants containing the inverted repeat (V line) showing very strong suppression. The majority (11 out of 13) of plants of the C line did not show suppression of ACO activity but overexpression, compared to wild-type plants, as would be expected since this construct contained a translatable ACO1 coding sequence.

To test for the presence of the transgenic ACO sequence, DNA from the plants was  
20 analysed by PCR using two oligos homologous to and complementary with the beginning and end respectively of the ACO1 coding sequence. This combination co-amplifies 1500 bp of the endogenous ACO1 gene (which acts as an internal positive control) and the ACO1 sense transgene as a 1000 bp fragment (since it was derived from a cDNA and so has no introns). The amplified region does not include the repeated region of the V-type transgene.  
25 The two fragments were separated by gel electrophoresis and detected by staining with ethidium bromide. This showed the presence of the transgene in all plants of the C line and all plants of the V line except one (V2) which also had no reduced ACC-oxidase activity (Figure 2).

- 11 -

**Example 1.2**

It was considered possible that the repeated region in the transgene might have affected the number of transgenes which integrated into the genome and that this was the actual source of high frequency silencing. The PCR assay described above can be used to estimate the transgene copy number if the following assumptions are made:

- 1) that in any transgenic plants there was no variation in the number of endogenous ACO1 genes per genome;
- 2) that the amplification efficiency ratio (endogenous ACO1 DNA: transgenic ACO1 DNA) is constant;
- 3) the reaction is sampled at low DNA concentration to minimise product re-annealing. Since we were only concerned with estimating the number of transgenes in the two lines relative to each other and not absolute quantification of transgene copy number, we did not employ synthetic combinations of "transgene" and "endogenous gene" DNA as standards.

After 20 cycles of amplification, gel-electrophoresis, Southern blotting, and hybridisation with a radioactively labelled ACO1 cDNA, the signal from endogenous and transgenic ACO1 DNA was visualised and quantified by phosphorimaging. The average transgene: endogenous gene ratio for the C line was 0.96 and for the V line 1.08 indicating that the repeat region in the V construct does not cause more T-DNAs to integrate during transformation.

**Example 1.3**

ACO1 mRNA increased in abundance following wounding and/or treatment of leaves with cycloheximide but accumulation was approximately five times greater after treatment with cycloheximide than after mechanical wounding which we have previously used as a stimulus. Wounding of cycloheximide treated leaves failed to elicit a further increase in ACO1 mRNA amount. We found cycloheximide to be a more reproducible inducer of ACO1 mRNA accumulation than mechanical wounding and so have used it in preference to

- 12 -

the latter in this study. No further increase in the abundance of ACO1 mRNA was observed when the concentration of cycloheximide was increased from 50 to 250 ug/ml (date not shown).

#### Example 1.4

5       The 5' end of ACO1 mRNA extracted from plants is heterogeneous but consists of two major species which differ by 2 bases. The 5' untranslated region (both the sense and duplicated antisense sequences) in both of the constructs (C and V) was made approximately 10 base pairs shorter than those of the endogenous gene. This allowed the discrimination of endogenous gene and transgene-derived transcripts by ribonuclease protection assays using a  
10       probe transcribed from a genomic ACO1 sequence which extended from the start of the 3' end of the 5' untranslated region to a *AccI* site, in the promoter of ACO1, 222 bases upstream. In RNA from wild type leaves, there were several bands which may arise from distinct RNA species or from breaking of RNA duplexes during digestion. Some of the bands seem more susceptible to the effects of antisense suppression than others (although the  
15       general trend is still suppression).

      In leaves from lines V4, V11 and V28 (all <10% ACO activity), there was extensive co-suppression of the endogenous transcripts (relative to wild-type) and the transgene transcripts (relative to those from a control transgene (line C1). V4, V11 and V28 all exhibited greater suppression than the homozygous ACO-antisense line (Hamilton et.al.  
20       Nature 346, 284-287(1990)).

      The use of the protein synthesis inhibitor cycloheximide as a stimulant of ACO1 RNA accumulation did not obviously alleviate the suppression of this RNA by the sense transgenes in lines V4, V11 or V28.

25       Although the endogenous genes transcript is unquestionably suppressed, it is possible that the inverted repeat within the 5' end of the V transgene transcript excludes the probe and causes the signal from the transgenic RNA to be underrepresented. This seems unlikely for the following reason. When a probe that was not excluded by the inverted repeat was used to analyse RNA from the V line, the mRNA signal (which, using this probe, is actually the sum of the endogenous and the transgenic RNAs) was still much less than in the wild type. The

- 13 -

data shows that in the absence of silencing, the abundance of the endogenous and transgenic RNAs are comparable.

#### Example 1.5

We chose to measure the abundance of unprocessed transcripts in total RNA extracts as a indirect measurement of transcription whilst simultaneously measuring the amount of processed mRNA. This was achieved using RNA probes transcribed from genomic sequences spanning introns in ribonuclease protection assays. Since the RNA analysed was from leaves frozen in liquid nitrogen and then extracted in strongly protein-denaturing conditions (phenol and detergent) there should have been little opportunity for any resetting of transcription during the process. There was a greater abundance of mRNA following treatment with cycloheximide although the total amount of mRNA in the ACO-AS plants was reduced. In the ACO-sense line, V11, there was little or no increase in the mRNA signal. It is likely that this mRNA signal is mainly from the transgene which is transcribed by the 35S promoter which is not cycloheximide inducible. In contrast, the abundance of the primary transcript in all RNA samples increased following cycloheximide treatment. This RNA species originates only from the endogenous ACO1 gene since the transgene has no introns. In all cases the suppressing transgene had little or no effect upon the abundance of the primary transcript.

#### Example 1.6

Cycloheximide strongly stimulated the accumulation of both the ACO1 primary transcript and mature mRNA. Quantification of the signal from primary transcripts and mature ACO1 RNA in wild type leaves before and after treatment with cycloheximide showed that there was a 6 fold increase in the abundance of unprocessed ACO1 RNA but a 13 fold increase in the amount of processed ACO1 RNA. The abundance of transgenic ACO1 RNA (transcribed from the 35S promoter) in the C line also rose upon treatment with cycloheximide.

#### Example 1.7

- 14 -

Two tandemly linked copies of the 5'UTR (each unit = 79bp; 74.7% (A+T)) were ligated in the inverted orientation between the CaMV 35S promoter and an almost full length ACO1 cDNA (Figure 1). Either unit of this direct repeat has the capacity to form a large cruciform structure with the 5'untranslated region immediately downstream. After  
5 *Agrobacterium*-mediated transformation with this construct, 26 out of 28 plants recovered from tissue culture exhibited suppressed ACO activity. A much lower frequency (2/15) of suppression was observed with a control construct which lacked the duplicated 5'UTR but was otherwise the same.

More transgenic plants were obtained with the V construct than with the control  
10 construct (as well as exhibiting the high HDGS frequency). It is likely that this is a direct result of reduced ethylene synthesis as a result of ACO gene suppression. Previous results have shown that greatly improved callus regeneration could be achieved after transformation with constructs which contained an ACO-antisense gene.

Of the two plants transformed with the repeat construct that showed no suppression,  
15 one, V2, may have had a truncated T-DNA or be an untransformed escape since the transgenic ACO1 sequence could not be amplified. Since the repeat contained DNA sequences already in the gene, it seems unlikely that it is this sequence per se which elicits the effect upon gene silencing. It is much more likely that it is the structure of the repeat DNA (or the transcribed RNA) which is the source of the high frequency of silencing  
20 observed. The repeat within the V construct was similar to that with the control construct

Most instances of HDGS are associated with complex transgenic loci that contain repeats or whole or part T-DNAs rather than simple single insertions but it is not known whether this is a primary determinant of suppression or an indirect effect. There are examples where apparently single transgenes are associated with gene silencing but these are  
25 in the minority and in at least some of these examples the T-DNAs contain internal repeats. The data presented here suggest that deliberate introduction of small repeats in a transgene can increase the number of transgenic lines in which homologous genes have been suppressed to almost 100%. Sense suppression could be obtained with the control construct but at a much lower frequency. The deliberate introduction of repetitive DNA into a  
30 transgene may substitute for a requirement for the insertion of repeated T-DNA units to



- 15 -

produce silencing. Although the PCR assay used here is not absolutely quantitative, it does suggest that the average transgene dosage is about 2 implying that some of the lines exhibiting suppression have single insertions. In several of our lines, the suppression obtained is profound (Figure 2) which makes this strategy even more attractive to those interested in specifically switching off gene expression. There is one previous report of the deliberate combination of repetitive DNA with a reporter gene effecting increased HDGS: Lohuis et al., Plant Journal, 8, 919-932 (1995) inserted a copy of a randomly isolated repetitive genomic sequence (RPS) upstream of GUS reporter gene and found that this element increased the frequency of variegation of transgene expression. This is an example of cis-inactivation, probably acts at the transcriptional level, and the authors considered it to be distinct from co-suppression/sense-suppression phenomena. Interestingly, the RPS element did not increase the frequency of complete silencing of the transgene. In our example, although the level of suppression is severe in many lines, it is not possible to say whether the degree of suppression is equal in all cells expressing the target gene or if the repeat has simply greatly increased the proportion of cells experiencing suppression.

### Example 1.8

#### Constructs and transformation

The tomato ACO1 cDNA, pTOM13 was released from its original cloning vector, pAT153, (Promega), creating pG31. pG31 was digested with EcoRI and the vector re-ligated to create pTRD. This removed the 5' end of the cDNA which contains approximately 90 base pairs of the 3'untranslated region in the antisense orientation at its 5' end which may have been introduced artefactually during the original cloning of the pTOM13 cDNA. The remaining ACO1 sequence was cut out from pTRD with EcoRI and HindIII and ligated into pT<sub>7</sub>.T<sub>3a</sub>18 (BRL) digested by EcoRI and the ends filled in with Klenow enzyme. The 5' untranslated region of the ACO1 transcript (minus approximately 10 bases at the 5' end) was amplified with Taq polymerase from oligo dT-primed cDNA of wounded tomato leaves with the primers 5' CATTCACTCTTCAATCTTTTG 3' (SEQ ID No.2) and 5' CTTAATTCTTGGTAAAGTGTTC 3' (SEQ ID NO.3). This DNA was rendered flush ended with T4 DNA polymerase and ligated with the filled in pTRF to create pMI1. This reconstituted the EcoRI site at the 5' end and yielded a translatable ACO1 cDNA

- 16 -

slightly shorter than the wild type ACO1 mRNA. Sequencing confirmed that the amplified ACO1 sequence was not mutated. pMI1 was digested with HindIII and partially with EcoRI and the fragment containing the ACO1 cDNA sequence was filled in with Klenow enzyme, and ligated with SmaI digested pDH51 to create pDHC1. This was digested with XbaI and HindIII, the filled in and the fragment containing the vector, 35S promoter and ACO1 cDNA religated to create pMI5. pMI7 contains two copies of the 5'UTR of ACO1 tandemly linked and inserted in the antisense orientation upstream of the 5'UTR of ACO1 in pMI5. This was made by amplifying the 5'UTR from tomato leaf cDNA (see above) with oligos 5' CATTCACTCTTCAATCTTTTG 3' (SEQ ID No.2) and 5'CTTAATTTCTTGGTAAAGTGTTTCC 3'(SEQ ID NO.3), polishing the DNA with T4 DNA pol and ligating it into a filled in Acc651 site in pMI5 upstream of the 5'UTR of the ACO1 sequence Acc651 (an isoshizomer of Kpn1 but which gives a 5' overhang). The construction was confirmed by sequencing.

pDHC1 and pMI7 were digested with BamHI, BglII and PvuII and the BamHI-PvuII fragments containing the CamV35S-ACO1cDNA sequences were cloned into Bin19 which had been cut by HindIII, filled in and then cut by BamHI. The resulting recombinants were called pBC1 and pBM17 respectively. These plasmids were transformed into *A. tumefaciens* LBA4404: and this used to transform tomato cotyledons (*Lycopersicon esculentum* var Ailsa Craig). Plants were regenerated from callus grown on 50µg.ml<sup>-1</sup> kanamycin.

## Example 1.9

### ACC-oxidase assays

ACC-oxidase activity was measured as the ability of plant tissue to convert exogenous 1-aminocyclopropane-1-carboxylic acid (ACC) to ethylene. Discs were cut from leaf lamina with a sharp cork borer and placed in contact with 0.5ml of 10mM NaH<sub>2</sub>PO<sub>4</sub>/Na<sub>2</sub>HPO<sub>4</sub> (pH7), and 10 mM ACC (Sigma) in 5 ml glass bottles which were then sealed with "Subaseal" vaccine caps (Fisons). After 1 hour at room temperature, the ethylene in the head space was measured by gas chromatography as described by Smith et al., 1986. Ethylene was also measured from bottles containing the solution but without leaf

- 17 -

tissue. These values were subtracted from the values obtained from the bottles containing leaf discs.

### Example 1.10

#### PCR analysis of transgenic plants

5 DNA was extracted from single leaves of wild type plants, plants homozygous for a ACO-antisense gene, and those transformed with the constructs of pBC1 and pBM17. Leaves were frozen in liquid nitrogen, briefly ground in eppendorf tubes with a disposable pipette tip, ground further after the addition of 200µl DNA extraction buffer (1% laurylsarcosine, 0.8% CTAB, 0.8M NaCl, 0.02M EDTA, 0.2M Tris/HCl (pH8)), heated to  
10 65°C for 15 minutes, extracted once with phenol/chloroform and the DNA precipitated from the aqueous phase by the addition of 0.6 volumes of isopropanol. The DNA was recovered by centrifugation, the pellets washed in 70% ethanol, dried and redissolved in 200ul of TE buffer. 1ul of this was used as template for simultaneous PCR amplification of the endogenous ACO1 gene and the transgene using the primers ACO1.1  
15 (ATGGAGAACTTCCCAATTATTAAGTTGGAAAAG SEQ ID NO 4) and the ACO1.2 (CTAAGCACTTGCAATTGGATCACTTTCCAT SEQ ID NO 5) for 21 cycles of 30 seconds at 95°C, 30 seconds at 65°C and 1 minute at 72°C. Amplified DNA was separated by electrophoresis in a 0.8% agarose/1xTBE gel and blotted onto HybondN+ in 0.4M NaOH for 6 hours. To detect the amplified ACO sequences, the DNA on the filter was hybridised  
20 with random prime labelled ACO1 cDNA. The filter was washed in 0.2xSSPE/1%SDS at 65°C followed by phosphorimaging of the radioactive signal.

### Example 1.11

#### Treatment of leaves with cycloheximide and mechanical wounding

Compound leaves were excised with a sharp scalpel blade and immediately placed  
25 under water solution of 50µl.ml<sup>-1</sup> cycloheximide (Sigma). Another 3 cm of the stalk was cut from the branch under the solution and the assembly was then left in a laminar airflow for six hours to allow the cycloheximide to enter the leaves.

- 18 -

To wound leaf tissue, individual leaflets were placed on a hard surface and diced with a sharp scalpel blade approximately 10 times transversely and 5 times longitudinally.

#### Example 1.12

##### Northern analysis of ACO mRNA in leaves treated with cycloheximide

5 RNA was extracted from cycloheximide treated leaves as follows. Tissue was frozen in liquid nitrogen and pulverised either in a coffee grinder (for fruit pericarp, see below) or in a mortar (for leaves). 5ml.gfw<sup>-1</sup> of RNA extraction buffer (Kirby's) was added and the frozen slurry ground further in disposable polypropylene centrifuge tubes with a glass rod. Once thawed, the mixture was extracted twice with phenol/chloroform and the nucleic acids  
10 precipitated by the addition of 2.5 volumes of ethanol, 1/10 volume 3M sodium acetate (pH5) and refrigeration at 20°C for 1 hour. After centrifugation at 3000xg for 10 minutes (40 minutes for a fruit extraction), the pellets were redissolved quickly in water (approximately 1ml per gram of tissue) and, an equal vol. of 2x DNA extraction buffer (1.4M NaCl, 2% CTAB, 100mM Tris/HCl (pH8)). Two volumes of precipitation buffer (1%CTAB, 50mM  
15 Tris/HCl (pH8)) were added to precipitate the nucleic acids (30 minutes at room temperature suffices) and the precipitate was collected by centrifugation (3000xg/15 minutes). This step was repeated except the pellets were dissolved in 1xDNA extraction buffer. After collection of the second precipitation, the pellets were redissolved in 0.5ml 1M NaCl and immediately reprecipitated with 2.5 volumes of ethanol (-20°C/30 minutes). After centrifugation  
20 (10000xg/10 minutes), the pellets were redissolved in 400µl water and extracted twice with phenol/chloroform. The nucleic acids were precipitated and collected as above redissolved in 400µl water. 46ul of 10 x One-Phor-All-Buffer (Pharmacia) was added with 50 units of RNAase-free DNAase (Promega) and the solutions incubated at 37°C for 30 minutes. They were extracted twice with phenol/chloroform, the RNA precipitated and collected as above  
25 and finally redissolved in 100-500ul of water. We have found that this relatively extensive purification is necessary if rare transcripts are to be detected by RPA. Also, the RNA re-dissolves readily which greatly reduces handling time when manipulating this RNA mixed with radioactive probe RNA.

- 19 -

50µg of leaf RNA was mixed with an equal volume of denaturation/loading solution (50% formamide; 25mM sodium phosphate (pH6.5); 10mM EDTA; 6.2% formaldehyde; 200µg.ml<sup>-1</sup> ethidium bromide) and separated by electrophoresis on a 25mM sodium phosphate (pH6.5) /3.7% formaldehyde /1.5% agarose gel in 10mM sodium phosphate (pH6.5)/3.7% formaldehyde with continuous buffer re-circulation. The separated RNA was blotted onto Genescreen (Dupont) hybridisation membrane in 10mM sodium phosphate (pH6.5). The autocrosslink setting on a Stratalinker (Stratagene) was used to covalently link the RNA to the filter. The filter was prehybridised and then hybridised with a 32P-random prime labelled ACO1 cDNA probe. The filter was washed in 0.2xSSPE/1%SDS at 65°C and then exposed to Kodak X-omat film between two intensifying screens at -70 for 24 hours. Subsequently the radioactivity in each band was measured by phosphorimaging.

### Example 1.13

#### Ribonuclease protection analysis

RNA was extracted from cycloheximide treated leaves and fruit described above.

15 RNA probes were transcribed with T7 RNA polymerase at 20°C with α-<sup>32</sup>P UTP (400Ci. mmol<sup>-1</sup>) as the sole source of UTP. After 1 hour incubation, RNAase-free DNAase was used to remove the template and the probe was further purified on 6%polyacrylamide/8M urea/1xTBE gels. The band containing the full length probe was visualised by autoradiography. The gel slice containing this RNA was excised and placed in 1ml probe elution buffer (0.5M ammonium acetate; 1mMEDTA; 0.2% SDS) for between 6 and 14 hours at 37°C. Typically, between 20m and 100µl of this would be co-precipitated with between 20 or 100µg of the RNA to be tested plus two yeast RNA controls. The precipitated RNAs were redissolved in 30µl hybridisation solution (80% formamide; 40mM PIPES/NaOH; 0.4M sodium acetate; 1mM EDTA pH should be 6.4) heated to 65°C for 10 minutes and hybridised at 42°C for between 2 to 14 hours. The longer hybridisation times were purely for convenience since we easily detected even rare transcripts after only 2 hours of hybridisation. 300µl of RNAase digestion buffer (5mM EDTA; 200mM sodium acetate; 10mM Tris/HCl. Final pH of solution should be 7.5) containing either RNAaseONE (Promega) or RNAase T1 (Ambion) was added to each tube except one containing yeast RNA which received RNAase digestion buffer without any ribonuclease. Incubation of the

- 20 -

digesting RNA was at either 25°C (RNAaseONE) or 37°C RNAaseT1) for 2-4 hours. RNAaseONE was inactivated by the addition of SDS to 0.5% and the protected, double stranded RNAs were precipitated with ethanol and sodium acetate. RNAaseT1 was inactivated and the double stranded RNAs were precipitated by the addition of the inactivation/precipitation solution provided with the RNAase protection kit from Ambion. The protected RNAs were redissolved in 5-10ul of denaturation/loading solution (80% formamide; 10mM EDTA; 0.1% bromophenol blue; 0.1% xylene cyanol; 0.1% SDS), heated to 95°C for 5 minutes and then separated by electrophoresis on a on 6-8% polyacrylamide/8M urea/1xTBE gels (the concentration of polyacrylamide depending on upon the sizes of the fragments to be separated). After electrophoresis, the gels were dried and exposed to Kodak x-omat film between two intensifying screens at -70 for the time indicated. The radioactivity was measured by phosphorimaging.

#### EXAMPLE 2.0

##### Construction of synthetic heterologous DNA inverted repeat.

A synthetic heterologous DNA invert repeat (SEQ ID No 11) was constructed by annealing two sets of synthetic oligos (HIR1 SEQ ID No 12 and HIR2 SEQ ID No 13 and HIR 3 SEQ ID No 14 and HIR 4 SEQ ID No 15) and ligating each set into pSK-(bluescript, Statagene) independently, to create pHIRA and pHIRB respectively. The invert repeat structure was created by digesting both pHIRA/B vectors with XhoI and NcoI and ligating the 42bp fragment from pHIRB into the pHIRA. The invert repeat structure was isolated from the pSK- vector using KpnI and cloned into the KpnI site immediately downstream of the CaMV35S promoter in the plant expression cassette pSIN to create pHIR-SIN.

The tomato ACO1 cDNA (pTOM13) coding sequence was amplified from its original cloning vector pAT153 (promega) using two oligonucleotide primers, 5' CTTTACCAAGAAGTGCACATGGAGAACTTCCC 3' SEQ ID No 6, and 5'GAATTGGGCCCTAAGCACTTGCAATTGG 3' SEQ ID No 7 which prime either side of the TOM13 coding sequence introducing ApaLI and ApaI sites respectively. The PCR product was digested with ApaLI and ApaI and the ends blunted in using Pfu polymerase

- 21 -

(Stratagene). The blunt PCR fragment was ligated into the SmaI site downstream of the invert repeat structure of pHIR-SIN to create pSIN-HIR-ACO.

The plant expression cassette from pHIR-ACO was isolated using AgeI and ligated into the binary vector pVB6 AgeI site to create pHIR-ACO SEQ ID No 10. The insert was orientated  
5 using restriction analysis to ensure that all the ORF that will be active in the plant were unidirectional. pHIR-ACO was transformed into *A.tumefaciens* LBA4404: and this used to transform tomato cotyledons (*Lycopersicum esculentum* var Ailsa Craig). Plants were regenerated from callus.

### Example 2.1

#### 10 Identification of Transgenic Plants

DNA was extracted from single leaves and extracted as described previously. Plants containing the HIR-ACO T-DNA insert were identified by PCR using an internal TOM13 sense primer (5' GCTGGACTCAAGTTTCAAGCCAAAG 3' SEQ ID No 8) and a NOS  
3'UTR (untranslated region) specific antisense primer  
15 (5'CCATCTCATAAATAACGTCATGC3' SEQ ID No 9)

### Example 2.2

#### ACC-oxidase assays

ACC-oxidase activity was measured as the ability of plant tissue to convert exogenous 1-aminocyclopropane-1-carboxylic acid (ACC) to ethylene. Small leaves were removed from  
20 shoots and wounded with a scalpel before being placed into a 2ml sealable vial, and left for 30minutes. The vials were then sealed and left for an hour at room temperature, after which the ethylene in the head space was measured by gas chromatography as described by Smith et al., 1986. Ethylene was also measured from wildtype, over-expressing (C12) and antisense down-regulated plant material.

- 22 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: ZENECA LIMITED  
(B) STREET: 15 STANHOPE GATE  
(C) CITY: LONDON  
(D) STATE: LONDON  
(E) COUNTRY: UNITED KINGDOM  
(F) POSTAL CODE (ZIP): W1Y 6LN  
(G) TELEPHONE: 01344 414521  
(H) TELEFAX: 01344 481112  
(I) TELEX: 847556

(ii) TITLE OF INVENTION: GENE SILENCING

(iii) NUMBER OF SEQUENCES: 15

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(v) CURRENT APPLICATION DATA:  
APPLICATION NUMBER:

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3681 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA"

## (vii) IMMEDIATE SOURCE:

(B) CLONE: SEQ ID NO 1

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

AAATTTGATA GATTCAGTTT TTATGTTTTT AGTGCTGATT ACAACATTGA AATTCTAAAT	60
TTAGAATTTA ATATTTATTA AATGTTAGTG CATTTATACA AATAACATAT TACATCTCAA	120
ATAATATTGA GTTTGTTAGA TTTTATTTGC CCTGATTCT TATCATAAAT AGGTTTTCT	180
TTTAGGAAAA GGTTTTGAAT TGAATTTCT TTTTGGTA GGAAAAAGT TAGGACTCTA	240
TAAATAGAGG CATGTCCTT CTAACCTAAT TAGCATTAC AATGTAGTT TAAGGGCTT	300
GAGAGTTTG GTTAGAGGA GAATTTGTGA ACCTCTCATG TATTCCGAGT GAATTGGTTG	360
AGGTTGTTT CCTCTGTATT TTGTACTCTC ATGTTTATAG TGGATTGCTC ATTTCTTTG	420
TGGACGTAGG TCGATTGACC GAACCACGTT AAATTTTGT GTCTTTTGGT ATATTTCTG	480
TTCTTCTTAC TCGTGGTCTT TCGAGTTTG CTTTGCTAGC TTCCGCGTT ACACCTGCTT	540



- 23 -

	ATTTTCGGTC	CTAACAAGTG	GTATCAGAGC	CAGATTCAAT	AATGGAGTCA	GGTGTAGTGG	600
	TTCGATAATC	GATGATTGAA	CCAAGTTAGA	AAGAGGTGTT	CATCTTGACG	GGTGTAGTTC	660
5	TAGCCGCAAC	CTTTTTGACA	GTAATGAAGA	TTTTGATGGA	GAAATTGTTT	CAGAGAGGTT	720
	CTCTGTGTTG	AGACATAAAT	TTTGTAAGG	AGATTATGGA	GAGGAGAAGC	AAGTTGTTGA	780
10	AGATTAAGTA	AAGAAGGTGG	ACAAATCTAT	TTTGTCAGAA	ATTCAGGCCA	AGGGGGAGAT	840
	TTGTTGGGTT	TTATTTGCCC	TGATTTTTTA	CCATAAATAG	GTTTTCCTTT	AAGGAAAAGG	900
	TTTTGAATTG	ACTATTCTTT	TTTTGGTAGG	AAAAGGTTTA	GGATTCTATA	AATAGAGGCA	960
15	TGTTCCCTTCT	AACTTAATTA	GCATTCACAA	TGTAGTTTTA	AGGGCTTTGA	GAGTTTTGGT	1020
	TAGAGGGAGA	ATTTGTGAAC	CTCTCATGTA	TTCCGAGTGA	ATTGGTTGAG	GTTGTTTCCC	1080
20	TCTGTATTTT	GTACTCTCAT	GTTTATAGTG	GATTGCTCAT	TTCCTTTGTG	GACGTAGGTC	1140
	GATTGACCGA	ACCACGTTAA	ATCTTTGTGT	CTTTGGTAT	ATTTCTCGTT	GTCTTCTTAC	1200
	TCGTGGTCTT	TCGAGGTTTG	CTTTGCTAGC	TTCCGCGTTT	ACACCTGCTT	ATTTGCGGTC	1260
25	CTAACAGAGT	TCGATGGGTT	GAATCTATAA	AAAGAAAAAT	ATACTCGTGA	TTCACGATTA	1320
	TTTATATGAA	AATATAATAA	ATATTGAATT	TCCTTTGCTA	TTTCTTATGT	TTACGTCTTT	1380
30	ATATTTCAAA	TTATTCCACC	AATACTGACA	AGCCCTAGGC	CATCTCTAGG	AAATTCATAC	1440
	AATTTTTTTT	TTGTTGTAA	CTAGTTAAAT	TGGCAGCCTT	AAAGATTATT	GTAAAATTCA	1500
	AGGCAACTTC	CTCAAGTACT	ACAACTACAT	TGTAACATCC	CAGTCAAAGT	GTCCTAAAAT	1560
35	TTTATAAAAT	TTGACACATG	AAACAATAGC	ACAATAAATT	TTAGTACTAT	TGCAGCCATG	1620
	GCCCATAGC	CATCATGTAT	TATAGTCAAA	ATGGGTCCTT	TTCCAATTG	TCTTGATCCC	1680
40	AAAATCCCTT	TGTAGGTAAG	ATGGTTCAAC	AAGGAACTAT	GACTCTTAAG	GTAGACTTGG	1740
	ACTCATAGAC	TTGTCATAAC	TCATAAAGAC	TTGGAATATA	ATAATTATTC	ATTTAAATTA	1800
	TAATTCTCTA	CTTTAATATC	TTCTACTATA	AATACCCTTT	CAAAGCCTCA	TTATTTGTAC	1860
45	ATCAAACATT	GATATTCATC	TCTTCAATCT	TTGTATTCA	CATATTCTAT	TTATTCAATA	1920
	CACTTAGGAA	AACACTTTAC	CAAGAAATTA	AGATGGAGAA	CTTCCCAATT	ATTAACCTGG	1980
50	AAAAGCTCAA	TGGAGATGAG	AGAGCCAACA	CCATGGAAAT	GATCAAAGAT	GCTTGTGAGA	2040
	ATTGGGGCTT	CTTTGAGGTA	ATCATAAATT	ACATAAACAT	ATTAATATGT	TTGTTTCAAT	2100
	TTATCAGTCA	TACTTTTCTC	TGTTTTAAAA	TTAATGTCAC	TTCAATATT	TAATAATTCG	2160
55	CATGACATGT	TTATAACACA	ACAAGATATA	GGTTACATTT	TGATACATTA	TATATAACTT	2220
	CTGTCACACG	ACTCAAAAGT	CTTCTTAAT	TTCTTGAATT	CAATGATCGA	TCAAACCTAAG	2280
60	ACACGTAAAA	TGAAACGGGG	AATAGTAATT	CTGTTTGCTT	ATGTGATCAT	TGTAGTTGGT	2340
	GAACCATGGA	ATTCCACATG	AAGTAATGGA	CACAGTAGAG	AAAATGACAA	AGGGACATTA	2400
	CAAGAAGTGC	ATGGAACAGA	GGTTTAAGGA	ACTAGTGGCA	AGTAAGGGAC	TTGAGGCTGT	2460

- 24 -

TCAAGCTGAG GTTACTGATT TAGATTGGGA AAGCACTTTC TTCTTGCGCC ATCTTCCTAC 2520  
TTCTAATATC TCTCAAGTAC CCGATCTTGA CGAAGAATAC AGGTACATAC ATGTGTCCTA 2580  
5 CATATTGCGT ATATAATAAA TAAACACAAA ATTTAAGTTA TATACGCTGA CAGTATAACT 2640  
AATTATAATG TTGTACCAAA TGATGCAGAG AGGTGATGAG AGATTTTGCT AAAAGATTGG 2700  
10 AGAAATTGGC TGAGGAGTTA CTTGACTTAC TCTGTGAAAA TCTTGGAATT GAAAAAGGTT 2760  
ACTTGAAAAA TGCCTTTTAT GGATCAAAAG GTCCCAACTT TGGTACTAAA GTTAGCAACT 2820  
ATCCACCATG TCCTAAGCCC GATTTGATCA AGGGACTCCG CGCTCATACA GACGCAGGAG 2880  
15 GCATCATACT TCTGTTCCAA GATGACAAAG TGAGTGGCCT TCAACTCCTC AAAGACGAGC 2940  
AATGGATCGA TGTTCTCTCC ATGCGCCACT CTATTGTGGT TAACCTTGGT GACCAACTTG 3000  
20 AGGTACAAGA TTCACTAAGT GTGTGTGTTT TTATCACTAT AACTTAGAAG TAGTAACTAA 3060  
AAATGGTATT AATGAAATGT TATAAAAACA GGTGATCACT AACGGGAAGT ACAAGAGTGT 3120  
GCTGCACAGA GTAATTGCAC AAACAGACGG GACACGAATG TCATTAGCCT CATTTTACAA 3180  
25 TCCAGGAAGT GATGCAGTAA TATATCCAGC AAAAAGTTTG GTTGAAAAAG AGGCAGAGGA 3240  
AAGTACACAA GTGTATCCAA AGTTTGTGTT TGATGATTAC ATGAAGTTAT ATGCTGGACT 3300  
30 CAAGTTTCAA GCCAAAGAGC CAAGATTGGA AGCAATGAAG GCAATGGAAA GTGATCCAAT 3360  
TGCAAGTGCT TAGATCCCAA TTCAATTAAA AAAATTGGTG TTTGAAAAAT ATATTTAAAT 3420  
ATAGCAATCT ATGTATACAC ATTATTGCT CTTCTTATGT ATGGTAGAAT AAAGTTAGTA 3480  
35 TTAAAAAGA TTGTGATTG CTGCATATGT ATCAAAAAGA GTCCTAATAT TTGTATCTAT 3540  
AAATAAGGTG CCTTCTAGTG AAATTATACA AATAATAATT TGGAGTGTAT TGTCTTTCT 3600  
40 CATGTAATTT AACTTTTAAG TATCTTACTT TACAATATAC TGTTCACTTA TTGAACATAT 3660  
TGAGTGATAT ATTGACTCAA T 3681

## (2) INFORMATION FOR SEQ ID NO: 2:

45

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 22 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

50

- (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PCR PRIMER"

55

- (vii) IMMEDIATE SOURCE:  
(B) CLONE: PCR PRIMER SEQ ID NO 2

60

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CATTCATCTC TTCAATCTTT TG

22

## (2) INFORMATION FOR SEQ ID NO: 3:

- 25 -

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 26 base pairs  
    (B) TYPE: nucleic acid  
5      (C) STRANDEDNESS: unknown  
    (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid  
    (A) DESCRIPTION: /desc = "PCR PRIMER"  
10

(vii) IMMEDIATE SOURCE:  
    (B) CLONE: PCR PRIMER SEQ ID NO 3  
15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:  
CTTAATTCT TGGTAAAGTG TTTTCC 26

20 (2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 33 base pairs  
    (B) TYPE: nucleic acid  
25      (C) STRANDEDNESS: unknown  
    (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid  
    (A) DESCRIPTION: /desc = "PCR PRIMER"  
30

(vii) IMMEDIATE SOURCE:  
    (B) CLONE: PCR PRIMER SEQ ID NO 4  
35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:  
ATGGAGAACT TCCCAATTAT TAACTGGAA AAG 33

40 (2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 30 base pairs  
    (B) TYPE: nucleic acid  
45      (C) STRANDEDNESS: unknown  
    (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid  
    (A) DESCRIPTION: /desc = "PCR PRIMER"  
50

(vii) IMMEDIATE SOURCE:  
    (B) CLONE: PCR PRIMER SEQ ID NO 5  
55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:  
CTAAGCACTT GCAATTGGAT CACTTCCAT 30

60 (2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 32 base pairs  
    (B) TYPE: nucleic acid

- 26 -

(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid  
5 (A) DESCRIPTION: /desc = "PCR PRIMER"

(vii) IMMEDIATE SOURCE:  
(B) CLONE: PCR PRIMER SEQ ID NO 6

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CTTTACCAAG AAGTGCACAT GGAGAACTTC CC 32

15 (2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:  
20 (A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid  
25 (A) DESCRIPTION: /desc = "PCR PRIMER"

(vii) IMMEDIATE SOURCE:  
(B) CLONE: PCR PRIMER SEQ ID NO 7

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GAATTGGGCC CTAAGCACTT GCAATTGG 28

35 (2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:  
40 (A) LENGTH: 25 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid  
45 (A) DESCRIPTION: /desc = "PCR PRIMER"

(vii) IMMEDIATE SOURCE:  
(B) CLONE: PCR PRIMER SEQ ID NO 8

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GCTGGACTCA AGTTTCAAGC CAAAG 25

55 (2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:  
60 (A) LENGTH: 23 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid

- 27 -

(A) DESCRIPTION: /desc = "PCR PRIMER"

(vii) IMMEDIATE SOURCE:

(B) CLONE: PCR PRIMER SEQ ID NO 9

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

10 CCATCTCATA AATAACGTCA TGC

23

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 1949 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

20 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA"

25 (vii) IMMEDIATE SOURCE:

(B) CLONE: PHIR-ACO SEQ ID NO 10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

30

ACCGGTGAAT TCCCATGGAG TCAAAGATTC AAATAGAGGA CCTAACAGAA CTCGCCGTAA 60

AGACTGGCGA ACAGTTCATA CAGAGTCTCT TACGACTCAA TGACAAGAAG AAAATCTTCG 120

35 TCAACATGGT GGAGCAGGAC ACGCTTGTCT ACTCCAAAAA TATCAAAGAT ACAGTCTCAG 180

AAGACCAAAG GGCAATTGAG ACTTTTCAAC AAAGGGTAAT ATCCGGAAAC CTCCTCGGAT 240

40 TCCATTGCCC AGCTATCTGT CACTTTATTG TGAAGATAGT GGAAAAGGAG GTGGCTCCTA 300

CAAATGCCAT CATTGCGATA AAGGAAAGGC CATCGTTGAA GATGCCTCTG CCGACAGTGG 360

TCCCAAAGAT GGACCCCCAC CCACGAGGAG CATCGTGGA AAAGAAGACG TTCCAACCAC 420

45 GTCTTCAAAG CAAGTGGATT GATGTGATAT CTCCACTGAC GTAAGGGATG ACGCACAATC 480

CCACTATCCT TCGCAAGACC CTTCTCTAT ATAAGGAAGT TCATTTTATT TGGAGAGGAC 540

50 AGGGTACCGC GGCACGGCCA GCCACGCCGC TGAGCCCGCA GTTCTCTGAG TTTCTGCGGG 600

CTCAGCGGCG TGGCTGGCCG TGCCGCCCAT GGGCGGCGGG GCTGCAGGAA TTCGATATCA 660

AGCTTATCGA TACCGTCGAC CTCGAGGGGG GGCCCGGTAC CGGATCCCCT GCACATGGAG 720

55 AACTTCCCAA TTATTAATT GGAAAAGCTC AATGGAGATG AGAGAGCCAA CACCATGGAA 780

ATGATCAAAG ATGCTTGTA GAATTGGGGG TTCTTTGAGT TGGTGAACCA TGAATTCCA 840

60 CATGAAGTAA TGGACACAGT AGAGAAAATG ACAAAGGGAC ATTACAAGAA GTGCATGGAA 900

CAGAGGTTTA AGGAACTAGT GGCAAGTAAG GGAATTGAGG CTGTTCAAGC TGAGGTTACT 960

GATTTAGATT GGGAAAGCAC TTTCTTCTTG CGCCATCTTC CTAATTCTAA TATCTCTCAA 1020

- 28 -

GTACCCGATC TTGACGAAGA ATACAGAGAG GTGATGAGAG ATTTTGCTAA AAGATTGGAG 1080  
 AAATTGGCTG AGGAGTTACT TGACTIONCTC TGTGAAAATC TTGGACTTGA AAAAGGTTAC 1140  
 5 TTGAAAAATG CCTTTTATGG ATCAAAAGGT CCCAACTTTG GTACTAAAGT TAGCAACTAT 1200  
 CCACCATGTC CTAAGCCCGA TTTGATCAAG GGACTCCGCG CTCATACAGA CGCAGGAGGC 1260  
 10 ATCATACTTC TGTTCOAAGA TGACAAAGTG AGTGGCCTTC AACTCCTCAA AGACGAGCAA 1320  
 TGGATCGATG TTCCTCCCAT GCGCCACTCT ATTGTGGTTA ACCTTGGTGA CCAACTTGAG 1380  
 GTGATCACTA ACGGGAAGTA CAAGAGTGTG CTGCACAGAG TAATTGCACA AACAGACGGG 1440  
 15 ACACGAATGT CATTAGCCTC ATTTTACAAT CCAGGAAGTG ATGCAGTAAT ATATCCAGCA 1500  
 AAAACTTTGG TTGAAAAAGA GGCAGAGGAA AGTACACAAG TGTATCCAAA GTTTGTGTTT 1560  
 GATGATTACA TGAAGTTATA TGCTGGACTC AAGTTTCAAG CCAAAGAGCC AAGATTTGAA 1620  
 20 GCAATGAAGG CAATGGAAAG TGATCCAATT GCAAGTGCTT AGGGGAGCCT GGGCCCCCTGC 1680  
 AGGTCGTTCA AACATTTGGC AATAAAGTTT CTTAAGATTG AATCCTGTTG CCGGTCTTGC 1740  
 25 GATGATTATC ATATAATTC TGTGAATTA CGAATTGCAT GTAATAATTA ACATGTAATG 1800  
 CATGACGTTA TTTATGAGAT GGGTTTTTAT GATTAGAGTC CCGCAATTAT ACATTTAATA 1860  
 CGCGATAGAA AACAAAATAT AGCGCGCAAA CTACCATAAA TTATCGCGCG CGGTGTCATC 1920  
 30 TATGTTACTA GATCGGGAAG CTTACCGGT 1949

## (2) INFORMATION FOR SEQ ID NO: 11:

- 35 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 78 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown  
 40  
 (ii) MOLECULE TYPE: other nucleic acid  
 (A) DESCRIPTION: /desc = "PCR PRIMER"  
 45  
 (vii) IMMEDIATE SOURCE:  
 (B) CLONE: SYNTHETIC INVERTED REPEAT SEQ ID NO 11

## 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GCGGCACGGC CAGCCACGCC GCTGAGCCCG CAGTTTCTCG AGTTTCTGCG GGCTCAGCGG 60  
 CGTGGCTGGC CGTGCCGC 78

## 55 (2) INFORMATION FOR SEQ ID NO: 12:

- 60 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 75 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown  
 (ii) MOLECULE TYPE: other nucleic acid

- 29 -

(A) DESCRIPTION: /desc = "PRIMER"

(vii) IMMEDIATE SOURCE:

5 (B) CLONE: PCR PRIMER SEQ ID NO 12

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

10 GCGGGTACCG CGGCACGGCC AGCCACGCCG CTGAGCCCGC AGTTTCTCGA GGATGGGTTG 60  
GCTCCATGGG CGGCG 75

15 (2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 75 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid

25 (A) DESCRIPTION: /desc = "PCR PRIMER"

(vii) IMMEDIATE SOURCE:

30 (B) CLONE: PCR PRIMER SEQ ID NO 13

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

35 CGCCGCCCAT GGAGCCAACC CATCCTCGAG AACTGCGGG CTCAGCGGCG TGGCTGGCCG 60  
TGCCGCGGTA CCCGC 75

(2) INFORMATION FOR SEQ ID NO: 14:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 66 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

45 (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "PCR PRIMER"

50 (vii) IMMEDIATE SOURCE:

(B) CLONE: PCR PRIMER SEQ ID NO 14

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GGGGCGCCGC TCGAGTTTCT GCGGGCTCAG CGGCGTGGCT GGCCGTGCCG CCCATGGCGC 60  
ATCGGG 66

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 66 base pairs

- 30 -

(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: unknown

5 (ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "PCR PRIMER"

(vii) IMMEDIATE SOURCE:  
10 (B) CLONE: PCR PRIMER SEQ ID NO 15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

15 CCCTAGGGCGC CATGGGCGGC ACGGCCAGCC ACGCCGCTGA GCCCGCAGAA ACTCGAGCGG 60

CGCCCC 66

20



- 31 -

## CLAIMS

1. A vector for enhancing the inhibition of a selected target gene within an organism,  
comprising a gene silencing vector characterised in that the said gene silencing vector  
5 includes a inverted repeat of all or part of a polynucleotide region within the vector.
2. A vector as claimed in claim 1, in which the inverted repeat sequence is a synthetic  
polynucleotide sequence and its inverted repeat sequence.
- 10 3. A vector as claimed in claim 1, in which the inverted repeat sequence is an inverted  
repeat of all or part of the said gene silencing vector.
4. A vector as claimed in claim 3, in which the inverted repeat sequence is an inverted  
repeat of the 5'-untranslated region of the gene silencing vector.
- 15 5. A method as claimed in any of claims 1 to 4, in which the inverted repeat is separated  
from the polynucleotide region by a sequence of nucleotides.
6. A method of controlling the expression of a DNA sequence in a target organism,  
20 comprising inserting into the genome of said organism an enhanced gene silencing  
vector as claimed in any of claims 1 to 4.
7. A vector for enhanced gene silencing comprising in sequence a promoter region, a 5'-  
untranslated region, a transcribable DNA sequence and a 3'-untranslated region  
25 containing a polyadenylation signal, characterised in that the said construct includes  
an inverted repeat of a region of said construct.
8. A vector as claimed in claim 7 in which the inverted repeat is a fragment of the 5'-  
untranslated region of the said construct.
- 30

- 32 -

9. A vector as claimed in claim 7 or claim 8, in which the inverted repeat is separated from the selected fragment by a sequence of nucleotides acting as a spacer.
10. A vector as claimed in claim 7 or 8 or 9, in which the construct includes a double  
5 copy of the inverted repeat.
11. A vector as claimed in any of claims 7 to 10, in which the vector two tandem copies of the inverted repeat.
- 10 12. A DNA construct for the inhibition of gene expression comprising in sequence a promoter region, a 5'-untranslated region, a transcribable DNA sequence and a 3'-untranslated region containing a polyadenylation signal, characterised in that the said 5'-untranslated region is contiguous with a pair of tandem inverted repeats of said 5'-untranslated region.
- 15

C

1/3

V

FIGURE 1A





FIGURE 1B



2/3

FIGURE 2

# Relative ACO activity in plants transformed with C and V constructs





FIGURE 3



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 98/01450

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C12N15/63 C12N15/82				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	DORER D. AND HENIKOFF S.: "Expansions of transgene repeats cause heterochromatin formation and gene silencing in <i>Drosophila</i> " CELL, vol. 77, no. 7, 1 July 1994, pages 993-1002, XP002075449 * see the whole document, esp. discussion, figure 5 *	1,3,7		
A	ASSAAD F. ET AL.: "Epigenetic repeat-induced gene silencing (RIGS) in <i>Arabidopsis</i> " PLANT MOLECULAR BIOLOGY, vol. 22, no. 6, September 1993, pages 1067-1085, XP002075450 * See the whole document, esp. p.1081 l. col. 1.30 - r. col. 1.3 * --- -/--	1-12		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.				
<input checked="" type="checkbox"/> Patent family members are listed in annex.				
<b>* Special categories of cited documents:</b>				
<table border="0"> <tr> <td style="vertical-align: top;">           "A" document defining the general state of the art which is not considered to be of particular relevance            "E" earlier document but published on or after the international filing date            "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)            "O" document referring to an oral disclosure, use, exhibition or other means            "P" document published prior to the international filing date but later than the priority date claimed         </td> <td style="vertical-align: top;">           "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention            "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone            "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art            "S" document member of the same patent family         </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "S" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "S" document member of the same patent family			
Date of the actual completion of the international search  28 August 1998		Date of mailing of the international search report  14/09/1998		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Kania, T		

Form PCT/ISA/210 (second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

National Application No  
PCT/GB 98/01450

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TEN LOHUIS M. ET AL.: "A repetitive DNA fragment carrying a hot spot for de novo DNA methylation enhances expression variegation in tobacco and petunia" PLANT JOURNAL, vol. 8, no. 6, December 1995, pages 919-932, XP002075451 cited in the application see the whole document	1-12
A	WO 97 01952 A (DNA PLANT TECHN CORP) 23 January 1997 * see esp. p.13-15 *	1-12
A	WO 93 23551 A (SEYMOUR GRAHAM BARRON ;TUCKER GREGORY ALAN (GB); GRIERSON DONALD ) 25 November 1993 see the whole document	1-12
A	GRIERSON, DON: "Silent genes and everlasting fruits and vegetables" NAT. BIOTECHNOL. (1996), 14(7), 828-829 CODEN: NABIF9;ISSN: 1087-0156, XP002075452 see the whole document	1-12
A	BLUME B ET AL: "Identification of transposon-like elements in non-coding regions of tomato ACC oxidase genes." MOLECULAR AND GENERAL GENETICS, (1997 APR 16) 254 (3) 297-303. JOURNAL CODE: NGP. ISSN: 0026-8925., XP002075453 see the whole document	1-12
A	HAMILTON, A. J. ET AL: "Post-transcriptional gene-silencing in tomato" MECH. APPL. GENE SILENCING, 'EASTER SCH. AGRIC. SCI.', 57TH (1996), MEETING DATE 1995, 105-117. EDITOR(S): GRIERSON, DONALD;LYCETT, GRANTLEY W.; TUCKER, GREGORY A. PUBLISHER: NOTTINGHAM UNIVERSITY PRESS, NOTTINGHAM, UK. CODEN: 63NBAT, XP002075454 see the whole document	1-12
T	STAM, M. ET AL: "Post-transcriptional silencing of chalcone synthase in Petunia by inverted transgene repeats" PLANT J., (19970700) VOL. 12, NO. 1, PP. 63-82. ISSN: 0960-7412., XP002075455 see the whole document	1-12

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

page 2 of 2

13:17:41

# INTERNATIONAL SEARCH REPORT

Information on patent family members

National Application No

PCT/GB 98/01450

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9701952 A	23-01-1997	AU 6402996 A EP 0837624 A	05-02-1997 29-04-1998
WO 9323551 A	25-11-1993	AU 4079493 A EP 0644942 A ZA 9303361 A	13-12-1993 29-03-1995 23-09-1994

Form PCT/ISA/210 (patent family annex) (July 1992)